

LRS Bianchi Type-V Anisotropic Dark Energy Cosmological Models With Time-Dependent Deceleration Parameter

Yogendra D. Patil

Department of Mathematics, Vidyan Mahavidyalaya, Malkapur(Dist. Buldana) 443101, India

ABSTRACT

Bianchi type-V cosmological models with variable equation of state (EoS) parameter have been investigated in general relativity when universe is filled with dark energy. The field equations have been solved by considering $R(t) = (t^n e^t)^{\frac{1}{l}}$ which gives $q = \frac{lr}{(t+r)^2} - 1$. This renders early decelerating and late time accelerating cosmological models. The physical and geometrical properties of the model are also discussed.

Keyword - Dark Energy, Deceleration Parameter, Bianchi Type-V Space Time.

1. INTRODUCTION

To study cosmological models, one of the important observational quantities is the deceleration parameter q . In any cosmological model, the Hubble constant H and deceleration parameter q play an important role in describing the nature of the evolution of the universe. The former one tells us the expansion rate of the universe today while the latter one characterizes the accelerating ($q < 0$) or decelerating ($q > 0$) nature of the universe. Some relativists assume various physical or mathematical conditions to obtain an exact solution of the Einstein's field equations. Berman [1,2] proposed a special law of variation for Hubble's parameter to obtain the cosmological solutions called the models with constant deceleration parameter (CDP) by assuming constant. This law is used by number of authors to study the cosmological models. Yadav [4] and Chawla et al. [5] have studied cosmological models with variable deceleration parameter.

In 2012, Saha et al. [14] have obtained cosmological solutions for FRW universe filled with two fluids consisting of dark energy and barotropic fluid by selecting the average scale factor which generates a time-dependent deceleration parameter such that the model generates a transition of the universe from early decelerating phase to the recent accelerating phase. Pradhan and Amirshachi [8] have also investigated accelerating dark energy models in Bianchi type-V space-time by selecting the scale factor as in Saha. However, Yadav [9,10], Pradhan [15], Rahman and Ansari [11] have generalized the average scale factor a given by , where n, m are positive constants and obtained the cosmological solutions. Recently Pradhan et al. [15] and Rahman et al. [11] have studied cosmological models based on Lyra's geometry with constant and time-dependent displacement field in different context.

Motivated by this study about the deceleration parameter from constant to time dependent, an attempt is made to study Bianchi Type-V space-time when the universe is filled with DE with time-dependent DP in general relativity. This work is organized as follows: In Section 2, the model and field equations have been presented. The field equations have been solved in Section 3 by choosing time depending deceleration parameters. The physical and geometrical behaviors of the models have been discussed in Sections 4. In the last Section 5, concluding remarks have been expressed.

2. METRIC AND FIELD EQUATION

Bianchi Type-V metric is considered in the form

$$ds^2 = dt^2 - a^2 dx^2 - b^2 e^{-2mx} (dy^2 + dz^2), \quad (1)$$

where $a(t)$ and $b(t)$ are scale factors and are functions of cosmic time t .

The energy-momentum tensor for a perfect fluid is

$$T_i^j = (\rho + p)u_i u^j - p g_i^j, \tag{2}$$

where p is the pressure, ρ is the energy density and g_j^i is a metric tensor. In co-moving coordinate system, u^i are the four co-moving velocity vectors which satisfy the condition

$$u^i u_i = 0, \quad \text{for } i = 1, 2, 3$$

and $u^i u_i = 1, \quad \text{for } i = 0.$

From equation (2), the components of energy-momentum tensor are

$$T_0^0 = \rho, \quad T_1^1 = T_2^2 = T_3^3 = -p. \tag{3}$$

With the help of equation (3), the matter tensor is given by

$$T_j^i = \text{diag}(\rho, -p, -p, -p). \tag{4}$$

For the perfect fluid, p and ρ are related by an equation of state

$$p = \omega \rho, \quad 0 \leq \omega \leq 1. \tag{5}$$

The Einstein's field equations are given by

$$R_i^j - \frac{1}{2} g_i^j R = -T_i^j, \tag{6}$$

where R_i^j is a Ricci tensor, R is the Ricci scalar.

The Ricci scalar for the Bianchi Type-V metric is given by

$$R = 2 \left(\frac{\ddot{a}}{a} + 2 \frac{\ddot{b}}{b} + \frac{\dot{b}^2}{b^2} + 2 \frac{\dot{a}\dot{b}}{ab} - 3 \frac{m^2}{a^2} \right).$$

With the help of equations (4) and (5), the field equations (6), for the metric (1) are

$$2 \frac{\dot{a}\dot{b}}{ab} + \frac{\dot{b}^2}{b^2} - 3 \frac{m^2}{a^2} = \rho, \tag{7}$$

$$2 \frac{\ddot{b}}{b} + \frac{\dot{b}^2}{b^2} - \frac{m^2}{a^2} = -\omega \rho, \tag{8}$$

$$\frac{\ddot{a}}{a} + \frac{\ddot{b}}{b} + \frac{\dot{a}\dot{b}}{ab} - \frac{m^2}{a^2} = -\omega \rho. \tag{9}$$

Here the over dot $\left(\dot{} \right)$ represents the differentiation with respect to t .

3. SOLUTION OF THE FIELD EQUATIONS

The field equations (7) to (9) are a system of three non-linear differential equations with four unknown parameters. The system is thus initially undetermined. To obtain a deterministic solution the following physical conditions are used.

(i) The expansion scalar (θ) is proportional to shear (σ). This condition leads to

$$\frac{1}{\sqrt{3}} \left(\frac{\dot{a}}{a} - \frac{\dot{b}}{b} \right) = \alpha_0 \left(\frac{\dot{a}}{a} + 2 \frac{\dot{b}}{b} \right),$$

which yields

$$\frac{\dot{a}}{b} = m \frac{\dot{b}}{b},$$

Where α_0 and m are arbitrary constants.

Above equation after integration reduces to

$$a = \eta(b)^m,$$

where η is an integration constant.

Here, for simplicity and without loss generality, we assume that $\eta = 1$.

Hence, we have

$$a = b^m, \quad m \neq 1. \tag{10}$$

Collins *et al.* have pointed out that for the spatially homogeneous metric the normal congruence to the homogenous expansion satisfies that the condition $\frac{\sigma}{\theta}$ is constant [12].

(ii) Now, extra condition is needed to solve the system completely. Hence, the model of deceleration parameters are considered as discussed in the following sections.

4. MODEL WITH TIME DEPENDENT DECELERATION PARAMETER

The law of variation of Hubble parameter proposed by Saha *et al.* [14] modified by Singha and Debnath [13] called as special form of deceleration parameter defined by

$$q = -\frac{R\ddot{R}}{\dot{R}^2} = \frac{lr}{(t+r)^2} - 1. \tag{11}$$

The proposed law yields a time-dependent DP which describes the transition of the universe from the early decelerating phase to current accelerating phase.

$$\begin{aligned} 1 - \frac{R\ddot{R}}{\dot{R}^2} &= \frac{lr}{(t+r)^2} \\ \Rightarrow \frac{\dot{R}^2 - \ddot{R}R}{\dot{R}^2} &= \frac{lr}{(t+r)^2} \\ \Rightarrow \frac{\partial}{\partial t} \left(\frac{R}{\dot{R}} \right) &= \frac{lr}{(t+r)^2}. \end{aligned}$$

After integration, the above equation simplifies to

$$\begin{aligned} \frac{R}{\dot{R}} &= \frac{lt}{t+r} \\ \Rightarrow \frac{\dot{R}}{R} &= \frac{t+r}{lt}. \end{aligned}$$

Again integrating, it reduces to

$$\begin{aligned} \log R &= \frac{1}{l} (r \log t + t) \\ &= \frac{1}{l} \log (t^r e^t) \\ \Rightarrow R &= (t^r e^t)^{\frac{1}{l}}. \end{aligned} \tag{12}$$

For the metric (1), the average scale factor R is given by

$$R = (ab^2)^{\frac{1}{3}}. \tag{13}$$

From the equations (12) and (13),

$$R = (ab^2)^{\frac{1}{3}} = (t^r e^t)^{\frac{1}{l}}$$

$$\Rightarrow ab^2 = (t^r e^t)^{\frac{3}{l}}.$$

Using equation(10), it reduces to

$$b^m b^2 = (t^r e^t)^{\frac{3}{l}}$$

$$\Rightarrow b = (t^r e^t)^{\frac{3}{(m+2)l}}. \tag{14}$$

With the help of equation(14), equation (10) leads to

$$a = (t^r e^t)^{\frac{3m}{(m+2)l}}. \tag{15}$$

Thus, the Bianchi Type-V anisotropic DE cosmological model in general relativity with time-dependent deceleration parameter can be written as

$$ds^2 = dt^2 - (t^r e^t)^{\frac{6m}{(m+2)l}} dx^2 - (t^r e^t)^{\frac{6}{(m+2)l}} e^{-2mx} (dy^2 + dz^2). \tag{16}$$

4.1 Physical properties of the model

For the cosmological model(16), the physical quantities such as spatial volume V , Hubble parameter H , expansion scalar θ , mean anisotropy parameter A_m , shear scalar σ^2 , energy density ρ and equation of state (EoS) parameter ω are obtained as follows:

The spatial volume is in the form

$$V = (t^r e^t)^{\frac{3}{l}}. \tag{17}$$

The rate of expansion H_i are

$$H_x = \frac{3m}{(m+2)l} (rt^{-1} + 1), \quad H_y = \frac{3}{(m+2)l} (rt^{-1} + 1). \tag{18}$$

The Hubble parameter is given by

$$H = \frac{1}{l} (rt^{-1} + 1). \tag{19}$$

The expansion scalar is

$$\theta = 3H = \frac{3}{l} (rt^{-1} + 1). \tag{20}$$

The mean anisotropy parameter is

$$A_m = \frac{2(m-1)^2}{(m+2)^2} = \text{constant} \neq 0, \text{ for } m \neq 1. \tag{21}$$

The shear scalar is given by

$$\sigma^2 = \frac{3(m-1)^2}{l^2(m+2)^2} (rt^{-1} + 1)^2. \tag{22}$$

It is observed that

$$\lim_{t \rightarrow \infty} \frac{\sigma^2}{\theta^2} = \frac{(m-1)^2}{3(m+2)^2} \neq 0. \tag{23}$$

The energy density is

$$\rho = \frac{9(2m+1)}{(m+2)^2 l^2} (rt^{-1} + 1)^2 - 3m^2 (t^r e^t)^{\frac{-6m}{(m+2)l}}. \tag{24}$$

The EoS parameter is

$$\omega = - \frac{\left[\frac{27}{(m+2)^2 l^2} (rt^{-1} + 1)^2 - \frac{6}{(m+2)l} rt^{-2} - m^2 (t^r e^t)^{\frac{-6m}{(m+2)l}} \right]}{\left[\frac{9(2m+1)}{(m+2)^2 l^2} (rt^{-1} + 1)^2 - 3m^2 (t^r e^t)^{\frac{-6m}{(m+2)l}} \right]} \quad (25)$$

For illustrative purposes, evolutionary behaviors of some cosmological parameters are shown graphically Fig -1.




Fig -1 Plots of time versus (a) Spatial volume (b) Energy density (c) Deceleration parameter for the values $r = 1, l = 1, m = 2$.

4.2 Physical behavior of the model

From Fig -1 (a), we observed that the spatial volume is zero at $t = 0$ and the expansion scalar is infinite showing that, the universe starts evolving with zero volume at $t = 0$ and expands with cosmic time t which is big bang scenario. Also, scale factors are zero at the initial epoch $t = 0$ hence the model has a point type singularity.

At $t \rightarrow \infty$, we have $q = -1$ and $\frac{dH}{dt} = 0$ indicating that the Hubble's parameter is maximum and the model has

the fastest rate of expansion for $t \rightarrow \infty$. The mean anisotropy parameter A_m is constant and $\lim_{t \rightarrow \infty} \frac{\sigma^2}{\theta^2} \neq 0 = const.$

is also constant, hence the model is anisotropic throughout the evolution of the universe except at $m = 1$ (i.e. the model does not approach isotropy). In Fig -1 (b), the plot of energy density versus time is given which indicates that the model starts with infinite density and as time increases the energy density tends to a finite value. Hence, after some finite time, the model approaches steady state. In Fig-1 (c), the plot of deceleration parameter versus time is given from which we conclude that the model is decelerating at an initial phase and changes from decelerating to accelerating. Hence the model is consistent with the recent cosmological observations [18-24]. Thus, our DE model is consistent with the results of recent observations.

5. CONCLUSIONS

A Bianchi Type-V cosmological model has been obtained when the universe is filled with DE in general relativity. To find a deterministic solution, we have considered the model of deceleration parameter which yields time-dependent scale factors.

In model (4), the solution of the field equations has obtained by choosing the time-dependent DP

$q = \frac{lr}{(t+r)^2} - 1$ which yields time-dependent scale factor $R = (t^r e^t)^{\frac{1}{l}}$. It is observed that the model has point type

singularity. In the early phase of the universe, the value of deceleration parameter is positive while as $t \rightarrow \infty$, the value of $q = -1$. Hence the universe had a decelerated expansion in the past and had accelerated expansion at present which is in good agreement with the recent observations of SN Ia.

It is worth to mention that in all cases, the models obtained are expanding, shearing, non-rotating and does not approach isotropy for large t . Further, the models are anisotropic throughout the evolution. Thus, DE models are in good harmony with recent cosmological observations [18-24]. We hope that the model will be useful for a better understanding of dark energy in cosmology to study an inflationary behavior of the Universe.

6. REFERENCES

1. Berman, M., *A special law of variation for Hubble's parameter*. Il Nuovo Cimento B (1971-1996), 1983. **74**(2): p. 182-186.
2. Berman, M.S. and F.D. Gomide, *Cosmological Models with Constant Deceleration Parameter*. General Relativity and Gravitation, 1988. **20**(2): p. 191-198.
3. Pradhan, A., J.P. Shahi, and C.B. Singh, *Cosmological models of universe with variable deceleration parameter in Lyra's manifold*. Brazilian Journal of Physics, 2006. **36**(4a): p. 1227-1231.
4. Yadav, A.K., *Some LRS Bianchi-I String Cosmological Models with Variable Deceleration Parameter*. arXiv preprint arXiv:1009.3867, 2010.
5. Chawla, C., R. Mishra, and A. Pradhan, *Anisotropic Bianchi-I cosmological models in string cosmology with variable deceleration parameter*. arXiv preprint arXiv:1203.4014, 2012.
6. Adhav, K., A. Bansod, and H. Ajmire, *Early decelerating & late time accelerating anisotropic cosmological models with dynamical EoS parameter*. Astrophysics and space science, 2013. **345**(2): p. 405-413.
7. Chirde, V. and S. Shekh, *Cosmological models with Magnetized Anisotropic Dark Energy in Lyra Geometry*. International Journal, 2014. **2**(6): p. 1103-1114.
8. Pradhan, A. and H. Amirhashchi, *Accelerating Dark Energy Models in Bianchi Type-V Spacetime*. Modern Physics Letters A, 2011. **26**(30): p. 2261-2275.
9. Yadav, A.K., *Bianchi-V string cosmological model and late time acceleration*. Research in Astronomy and Astrophysics, 2012. **12**(11): p. 1467-1474.
10. Kumar, Y.A., *Cosmological constant dominated transit universe from the early deceleration phase to the current acceleration phase in Bianchi-V spacetime*. Chinese Physics Letters, 2012. **29**(7): p. 079801.
11. Rahman, M.A. and M. Ansari, *Anisotropic Bianchi Type-III Dark Energy Model with Time-Dependent Deceleration Parameter in Sáez-Ballester Theory*. IOSR Journal of Applied Physics, 2013. **4**: p. 79-84.
12. Collins, C.B., E.N. Glass, and D.A. Wilkinson, *Exact Spatially Homogeneous Cosmologies*. General Relativity and Gravitation, 1980. **12**(10): p. 805-823.
13. Singha, A.K. and U. Debnath, *Accelerating Universe with a Special Form of Decelerating Parameter*. International Journal of Theoretical Physics, 2009. **48**(2): p. 351-356.
14. Saha, B., H. Amirhashchi, and A. Pradhan, *Two-fluid scenario for dark energy models in an FRW universe-revisited*. Astrophysics and Space Science, 2012. **342**(1): p. 257-267.

15. Pradhan, A., A.K. Singh, and D.S. Chouhan, *Accelerating Bianchi Type-V Cosmology with Perfect Fluid and Heat Flow in Saez-Ballester Theory*. International Journal of Theoretical Physics, 2013. **52**(1): p. 266-278.
16. Rao, V., et al., *Two-Fluid Scenario for Higher Dimensional Dark Energy Cosmological Model in Saez-Ballester Theory of Gravitation*. Prespacetime Journal, 2016. **7**(2).
17. Adhav, K.S., R.P. Wankhade, and A.S. Bansod, *LRS Bianchi type-I cosmological model with anisotropic dark energy and special form of deceleration parameter*. Journal of Modern Physics, 2013. **4**(08): p. 1037.
18. Perlmutter, S., et al., *Discovery of a supernova explosion at half the age of the Universe*. Nature, 1998. **391**(6662): p. 51-54.
19. Perlmutter, S., et al., *Measurements of Omega and Lambda from 42 high-redshift supernovae*. Astrophysical Journal, 1999. **517**(2): p. 565-586.
20. Perlmutter, S., et al., *Measurements* of the Cosmological Parameters Ω and Λ from the First Seven Supernovae at $z \geq 0.35$* . The astrophysical journal, 1997. **483**(2): p. 565.
21. Riess, A.G., et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*. Astronomical Journal, 1998. **116**(3): p. 1009-1038.
22. Riess, A.G., et al., *Type Ia supernova discoveries at $z > 1$ from the Hubble Space Telescope: Evidence for past deceleration and constraints on dark energy evolution*. Astrophysical Journal, 2004. **607**(2): p. 665-687.
23. Schmidt, B.P., et al., *The High-Z Supernova Search: Measuring cosmic deceleration and global curvature of the universe using Type IA supernovae*. Astrophysical Journal, 1998. **507**(1): p. 46-63.
24. Garnavich, P.M., et al., *Supernova limits on the cosmic equation of state*. Astrophysical Journal, 1998. **509**(1): p. 74-79.
25. MacCallum, M.A.H. (1971) A Class of Homogeneous Cosmological Models III: Asymptotic Behaviour. Communications in Mathematical Physics, 20, 57-84. <http://dx.doi.org/10.1007/BF01646733>